



### SICOBIO® RANGE

Product code: M-NPK333HO HS Code: 3101 00 00

# SICOBIO® NPK 3.3.3 (30% OM) Liquid High Organic fertiliser

- Made in E.U. -

04/2021

#### 1. PRODUCT DESCRIPTION & PROPERTIES

SICOBIO NPK 3.3.3 Liquid High Organic is best used as a base fertiliser, supplemented by SICOBIO NITRO and SICOBIO PHOS 8% to your crop's specific nitrogen and phosphate need. Due to the special production process, the nutrients are directly available to the plant. This product with high organic matter content is rich in fulvic and humic acids. These acids bind to nutrients in the soil, improving the roots' nutrient uptake. The product is fully made from plant-based materials and registered for use in organic agri- and horticulture. Due to the natural origin the nutrient content may vary.

## 2. PRODUCT SPECIFICATIONS & CERTIFICATE OF ANALYSIS Chemical Analysis

	% w/w
Total Nitrogen (N)	3 %
Organic Nitrogen	1.8 %
Ammoniacal Nitrogen (NH4-H)	1.2 %
Phosphorus (P2O5)	3 %
Potassium (K2O)	3 %
Sulphur (SO3)	3.1 %
Fulvic acids	16.3 %
Organic Matter	30 %
pH	4 - 6

#### **Physical Properties**

Appearance	Liquid
Colour	Light brown
Density (kg/l)	1.14 – 1.18

#### **3 ADVANTAGES & BENEFITS**

- Increases resistance against stress conditions
- · Prevents mineral deficiency
- Better rooting system
- · Higher cellular division
- Better metabolic and enzymatic activity
- · Improves soil properties
- Improves soil structure
- Better nutrient uptake

### 4. USAGE & RECOMMENDATIONS

#### \* Application

It is advised to always have an expert make a fertilisation plan and to start with a test application. No guarantees can be given in the event of incorrect use.

General recommended guidelines

APPLICATION		DOSE	DOSAGE
Foliar		3 – 5 ml/litre	Every 1 – 2 weeks
Irrigation	Tree nursery	30 – 50 litre/ha	Every 2 weeks
	Fruits and horticulture	0.4 – 0.8 % of irrigation water	Every week





Crop specific applications

CROP	APPLICATION
Vegetables	Start for 5 to 6 weeks during development of the first roots. Use weekly during vegetative development, fruit formation and hydric stress.
Cotton	First 5 weeks after planting. Weekly applications from just before bloom until the last bud has developed.
Corn	Start for 5 to 6 weeks during development of the first roots. 3-4 applications every 7-15 days from bloom until the last bud has developed.
Citrus and fruit trees	Start when the first leaves or buds form. Repeat every 7-15 days.
Grapevine and olive	Every 7-15 days from bloom until fruit development.
Ornamental	Start for 5-6 weeks during development of the first roots. Apply weekly, especially during bud formation.
Potato	During development of tubers, especially in sandy soils and at the end of the vegetative stage.

**<u>5. PACKING</u>** 0.25 − 0.5 − 1 − 10 − 20 -220 -1000 ltr

<u>6. STORAGE & SHELF LIFE</u> Store in a dark and dry environment at 10- 25° C.

Shelf life: 1.5 – 2 years if stored properly.