



**PRODUCT INFO
& DATASHEET**

Foliar Application Table

Foliar application timings (600 - 800 g/ha, 9 - 12 ounces/acre)

VEGETABLES	First	Second	Third	Optional
Eggplants, Pepper, Melon, Squash later	Six leaf stage	Early bloom	First fruit set	Four weeks
Bean, Peas	Six leaf stage	First bloom	First pods	
Cabbage, Broccoli, Cauliflower	Six true leaf stage	Three weeks later	Head initiation	
Cucurbits, Cucumbers	Six leaf stage	Just prior to bloom	Whilst picking	
Maize, Sweet Corn	Six leaf stage	55 - 75 cm / 20 - 30 inch growth	Just prior to tasseling	
Onions, Carrots, Leeks, Turnips	2 - 3 weeks after emergence	Root enlargement	Every 2 weeks until harvest	
Potatoes	Six leaf stage	When tuber approx. pea size	Early bloom	Bulking up
Tomato (fresh fruit market)	Six leaf stage	Early bloom	Whilst picking	After 2 weeks

FRUIT	First	Second	Third	Fourth	Optional
Grapes	20 - 30 cm cane	45 - 60 cm cane	Full blooming date	Berry set / early shattering	3 weeks later
Apples, Pears	Green tip	Pre-bloom pink bud	Full blooming date	Early fruit formation	Every 3 weeks
Citrus	Early bloom	Petal fall	With summer spray	With fall spray	Every 3 weeks
Bananas	Plant emergence	Adequate leaf size	Fruit half size	3 weeks later	Every 3 weeks
Strawberry and other berries	Just after transplant	Prior to bloom	Whilst picking	3 weeks later	Every 3 weeks
Stone fruit	Bud birst	Petal fall	4 weeks later	3 weeks later	Every 3 week
Plums, Cherries	Pink white bud	Full bloom	Early fruit formation	3 weeks later	Every 3 weeks

Foliar application timings (400 g/ha, 5 - 6 ounces/acre)

CEREALS	First	Second	Optional
WinterWheat and Oats	First node	Flag leaf	After any environmental stress
Winter (Malting) Barley	Early post emergence		After any environmental stress
Spring Wheat and Oats	1 - 3 Tillers	First node	Flag leaf and after any environmental stress
Spring (Malting) Barley	1 - 3 Tillers	First node	Flag leaf and after any environmental stress
Corn	At 2 - 6 leaf stage	At 50 - 75 cm growth	Just prior to tasseling

Foliar application timings (600 - 800 g/ha, 9 - 12 ounces/acre)

ECONOMIC CROP	First	Second	Third	Optional
Cotton	Seedling stage	Stage of flower emergence	Fully bloom	Every 4 weeks
Oilseed crops	Before reproductive growth stage	At height of flowering		Every 4 weeks
Tobacco	Early post emergence	After every leaves picked	Early bloom	Every 4 weeks
Hops	Early post emergence	Five weeks later		Every 4 weeks
Luceme		After each cut		

Plant

Turf (600 - 800 g/ha, 9 - 12 ounces/acre)
 year.
 Golf courses
 Greens (600 - 800 g/ha, 9 - 12 ounces/acre)
 Fairways (600 - 800 g/ha, 9 - 12 ounces/acre)

Foliar application

- 1) Recommended to apply "little and often", once or twice a month throughout the year.
- 2) Regular applications to greens and tees are especially beneficial.
- 3) Applied as bio-stimulant to increase the number and activity of micro-organism. This will improve grass growth and help reduce soil borne disease symptoms in the turf.
- 4) Tillering will also be increased, without excessive top growth.

Any information in this publication is believed to be accurate and is given in good faith, but is for the customer to satisfy itself of the suitability for its own particular purpose. No representation, warranty or guarantee is made to its accuracy, reliability or completeness.

SAP INTERNATIONAL CORPORATION by Krekelenberg 83, B-2980 Zoersel, Belgium
 Tel. +32-3-309.06.51 Email : info@sico.be Website : www.sico.be