





Nutritional Oral Liquids

SICOLYTES OS

Electrolytes for oral administration

1. Composition:

| Contains per ml: | |
|-----------------------------|----------|
| Dextrose | 280 mg. |
| Sodium chloride | 110 mg. |
| Glycine | 45 mg. |
| Sodium dihydrogen phosphate | 22 mg. |
| Potassium chloride | 13.5 mg. |
| Sodium citrate | 5 mg. |
| Solvents ad. | 1 ml. |

2. Description

Solution for oral administration.

The electrolytes sodium, phosphate, citrate, chloride, potassium and dextrose can be used for recovery of electrolyte and acid/base imbalances. The essential amino-acid glycine is added for a quicker recovery of the dehydrated animals.

3. Indications:

Prevention and treatment of dehydration caused by diarrhoea in calves, cattle, goats, poultry, sheep and swine.

4. Side effects:

No undesirable effects are to be expected when the prescribed dosage regimen is followed.

5. Dosage:

For oral administration through drinking water.

| Cattle, calves, goats and sheep | : | 15 - 30 ml per animal for 7 days. (Dilute 15 ml per 1.5 litre drinking water) |
|---------------------------------|---|---|
| Rabbits | : | 1 ml per animal for 7 days. (Dilute 1 ml per 100 ml drinking water.) |
| Poultry and swine | : | 1 litre per 2000 litres drinking water for 5 days. |

6. Withdrawal times:

None.

7. Warning:

Keep out of reach of children.

8.Packing:

Bottle containing 100, 500 and 1000 ml.