



**PRODUCT INFO  
& DATASHEET**

Nutritional Oral Liquids

# SICOMIN Cu-Chel Oral

## Solution for oral administration

### **1. Composition:**

Contains per ml:

Copper : 100 mg

### **2. Description**

SICOMIN Cu-Chel Oral is a nutritional oral liquid containing chelated copper. Chelated copper is easy to absorb and can be simply administered through drinking water. SICOMIN Cu-Chel Oral is mainly used to correct copper deficiency. Copper has beneficial effects on animal health and is a cofactor for many enzyme systems. Furthermore, copper can reduce the chance of infections, improves feed conversion and stimulates growth.

### **3. Benefits**

- complements copper deficiency
- chelated copper can easily be absorbed
- copper reduces the chance of infections
- copper is essential for the function of many enzyme systems
- improves feed conversion
- stimulates growth
- simple administration through drinking water

### **4. Indications**

SICOMIN cu-Chel Oral is indicated for the prevention and treatment of copper deficiency (hypocuprosis) and clinical conditions associated with copper deficiency.

Contra indications : - administration to ovine species and calves  
- administration to animals with known hypersensitivity to cupric substances  
- administration to animals with a seriously impaired hepatic and/or renal function  
- concurrent administration of other forms of copper treatment or supplementation

Side effects : overdosage may lead to copper toxicity

Withdrawal times : none

### **5. Dosage**

For oral administration via drinking water.

- Pigs and poultry: 100 ml SICOMIN Cu-Chel Oral per 1000 – 4000 litres of drinking water (2.5 – 10 ppm).

Note: The dosage and frequency of therapy required depends on the clinical condition and copper status of the animal as assessed by blood and liver levels both before and after therapy.

Only for use when there is a confirmed copper deficiency. High dietary levels of Molybdenum and Sulphur reduce Copper bioavailability.

### **6. Warning**

- keep out of the reach of children
- if there is any doubt about the copper status, random blood and/or liver samples should be taken prior to treatment
- high dietary levels of Mo and S reduce Cu bioavailability.
- after contact with eyes wash with plenty of water and seek medical advice
- wear suitable gloves and clothing
- do not mix SICOMIN Cu-Chel Oral with other medicines

### **7. Packaging**

Bottle of 1000 ml and jerrycan of 5 litres.

Any information in this publication is believed to be accurate and is given in good faith, but is for the customer to satisfy itself of the suitability for its own particular purpose. No representation, warranty or guarantee is made to its accuracy, reliability or completeness.

**SAP INTERNATIONAL CORPORATION** bv Krekelenberg 83, B-2980 Zoersel, Belgium  
Tel. +32-3-309.06.51 Email : [info@sico.be](mailto:info@sico.be) Website : [www.sico.be](http://www.sico.be)